

Scanning SQUID Microscope Imaging of Excess Vortex Density in Field-cooled MoGe Films with Surface Trenches

Dale J. Van Harlingen


DMR-0107253



University of Illinois at Urbana-Champaign


Motivation: Determine the effect of surface steps and defects on the distribution and dynamics of magnetic vortices in weak-pinning superconducting films

Approach: Scanning SQUID Microscope (SSM) images of the vortex density of amorphous-MoGe thin films patterned into squares and with trenched ion-milled into surface to simulate surface steps



Results:

- Vortex pattern with six-fold orientation but no long-range order even at low fields ($<0.1\text{G}$)
- Repulsion of vortices from trench edge
- Alignment of vortices on inside trench edge
- Excess net vortices in entire sample compared to applied field → **paramagnetic Meissner effect**





SSM images of vortex distributions in MoGe films without and with surface trenches at $B=30\text{mG}$

Scanning SQUID Microscope Imaging of Excess Vortex Density (cont'd)

D. J. Van Harlingen --- University of Illinois at Urbana-Champaign --- DMR-0107253

Experiments:

- Density of vortices inside trench is enhanced compared to the applied field density B/Φ_0 --- enhancement is independent of trench depth
- Excess vortices in trench exceeds depletion outside the trench edges \rightarrow net excess of vortices relative to expected density B/Φ_0



Conclusions: anomalous vortex density enhancement, a new paramagnetic Meissner effect ... origin may be screening currents or Type-III superconductivity along step edges

Personnel:

Adele Ruosi (Naples) --- visiting scientist
Britton Plourde --- graduate research student
(Ph. D. 2000: present address = UC Berkeley)
Micah Stoutimore --- graduate research student
Martin Stehno --- undergraduate student

Collaborators:

Peter Kes, Rut Besseling --- Leiden
Maurizio Russo, Carmine Granata
--- Naples, INFM